

Approximate solutions of the fractional Harry DYM equation

Sevil ÇULHA ÜNAL¹

ABSTRACT

In this paper, the approximate solutions of the time fractional Harry Dym equation with fractional derivative in the Caputo sense are obtained by using the Residual power series method (RPSM). This equation is a significant dynamical equation that occurs in a variety of physical systems. The suggested method provides good accuracy for the approximate solution when compared numerically with the exact solution. The effectiveness of the proposed method is also illustrated with the aid of numerical results. These results indicate that the RPSM is a power, useful, and applicable for determining the solutions of the time Harry Dym equation. Some of these results are illustrated by 2D and 3D graphics. Besides, the proposed method can be applied to many different differential equations due to its ease of use and reliability.

Keywords: *Approximate solution; Fractional partial differential equation; Caputo derivative; Harry Dym equation; Residual power series method.*

INTRODUCTION

The Harry Dym equation is in the form

$$\frac{\partial u}{\partial t} = u^3 \frac{\partial^3 u}{\partial x^3}$$

was first studied by Kruskal and Moser and is referred to an unpublished work of Harry Dym. This equation is entirely integrable nonlinear evolution equation linked to the traditional string problems [1]. More detailed information about these problems can be seen in [2-5]. The Harry-Dym equation is also closely related to the Korteweg-de Vries equation [6]. In the literature, numerous methods have been utilized to solve this equation. The solution methods for the Harry Dym equation are moving frame [7], Adomian decomposition [8], He's variational iteration [8], direct integration [8], power series [8], residual power series [8], Bäcklund transform [9], new iterative method [10], Haar wavelet [11], homotopy perturbation [12], reconstruction of variational iteration [12], Darboux transformation [13], and nonlinear steepest descent [14].

Recently, it has become very popular for scientists to obtain solutions of the fractional differential equations. These equations are widely used to model problems in viscoelasticity, turbulence, electrical networks, nonlinear biological systems, control theory, thermodynamics, fluid dynamics, signal processing, and so on [15-20]. The time fractional Harry Dym equation is one of the most important of them. So far, many researchers have used various analytical and numerical methods to obtain the time fractional Harry Dym equation. These methods are Adomian decomposition [21,22], homotopy perturbation Sumudu transform [22], Elzaki transform technique [23], Lie symmetry group analysis [24], similarity [25,26], homotopy analysis [27,28], Lie classical [29], homotopy perturbation [30], Mohand homotopy perturbation transform scheme [31], reduced differential transform [32], finite difference [33], q-homotopy analysis [34], and optimal system [35]. However, it is seen that the time fractional Harry Dym equation has not yet been solved with the RPSM.

The RPSM, proposed by Abu Arqub in 2013, is an efficient approach to obtain the approximate solutions of the different differential equations. These solutions are gained without the need for linearization, discretization, or perturbation. The RPSM does not require comparing the coefficients of the corresponding terms and does not

¹ Suleyman Demirel University, Department of Avionics, Isparta, Turkey

E-mail address: sevilunal@sdu.edu.tr

Orcid id: <https://orcid.org/0000-0001-7447-9219>

Manuscript Received 19 February 2023, Accepted 13 May 2023

This paper was recommended for publication in revised form by Regional Editor Ahmet Selim Dalkilic

¹ Suleyman Demirel University, Department of Avionics, Isparta, Turkey

* E-mail address: sevilunal@sdu.edu.tr

Orcid id: <https://orcid.org/0000-0001-7447-9219> Sevil Çulha Ünal

Manuscript Received 19 February 2023, Revised 18 April 2023, Accepted 13 May 2023

need a recursion relation. By selecting an appropriate value for the initial guesses approximations, the proposed method can be also directly applied to the equations. Besides, with this method, high precision is achieved by utilizing less time and small calculations. Moreover, by minimizing the residual error, the suggested method provides an easy way to achieve the convergence of the series solution. Furthermore, the RPSM relies on derivation, which is more accurate and much easier than integration. This is the basis of most other solution methods. In addition to all these, the proposed method suggests obtaining infinite series solutions with iterated operations.

In the present paper, the RPSM is used to get the approximate solutions of the time fractional Harry Dym equation of the form

$$D_t^\beta u(x, t) = u^3(x, t)u_{xxx}(x, t), \quad 0 < \beta \leq 1 \quad (1)$$

by the initial condition

$$u(x, 0) = \left(4 - \frac{3}{2}x\right)^{\frac{2}{3}} \quad (2)$$

where $D_t^\beta u$ is the Caputo fractional derivative of order β with respect to the time variable t . When $\beta = 1$, Eq. (1) turns into the standard Harry Dym equation. The exact solution for the Harry Dym is

$$u(x, t) = \left(4 - \frac{3}{2}(x + t)\right)^{\frac{2}{3}}.$$

The plan of this paper is as follows. In Section 2, the definitions and theorems of the Caputo derivative and the fractional power series are mentioned. In Section 3, the basic idea of the RPSM is expressed. In Section 4, the RPS solutions for the time fractional Harry Dym equation are obtained by suggested method. Besides, the efficiency and the reliability of this method are demonstrated by table and figures. In Section 5, the Conclusions are given.

PRELIMINARIES

There are numerous definitions of fractional operators, such as Grunwald-Letnikov, Caputo, Riemann-Liouville, Hadamard, Wely, and Marchaud in the literature. In this part, Caputo's definition is utilized since the derivative of a constant is zero and the initial conditions for the fractional differential equations with Caputo derivative take the familiar manner of integer order differential equations. The definition of Caputo derivative is defined as follows:

Definition 1. [36] The time fractional derivative of $u(x, t)$ in Caputo form is described as

$$D_t^\beta u(x, t) = \begin{cases} \frac{1}{\Gamma(m - \beta)} \int_0^t (t - \tau)^{m-1-\beta} \frac{\partial^m u(x, \tau)}{\partial \tau^m} d\tau, & m - 1 < \beta < m \\ \frac{\partial^m u(x, t)}{\partial t^m}, & m = \beta \in \mathbb{N}. \end{cases}$$

The definition and theorems for the fractional power series are given below. Details of them can be found in [37].

Definition 2. [37] A power series expansion of the manner

$$\sum_{m=0}^{\infty} c_m (t - t_0)^{m\beta} = c_0 + c_1 (t - t_0)^\beta + c_2 (t - t_0)^{2\beta} + \dots, \quad 0 \leq m - 1 < \beta \leq m, \quad t \geq t_0,$$

is called the fractional power series about t_0 . Here, t is a variable and the c_m 's are constants.

Theorem 1. [37] Assume that g is a fractional power series representation at t_0 of the manner

$$g(t) = \sum_{m=0}^{\infty} c_m (t - t_0)^{m\beta}, \quad 0 \leq m - 1 < \beta \leq m, \quad t_0 \leq t < t_0 + R.$$

If $D^{m\beta} g(t)$ are continuous on $(t_0, t_0 + R)$, then coefficients c_m are expressed as

$$c_m = \frac{D^{m\beta} g(t_0)}{\Gamma(m\beta + 1)}, \quad m = 0, 1, 2, \dots,$$

where R is the radius of convergence and $D^{m\beta} = D^\beta \cdot D^\beta \dots D^\beta$.

Theorem 2. [37] Assume that $u(x, t)$ has a multiple fractional power series representation at t_0 of the manner

$$u(x, t) = \sum_{m=0}^{\infty} g_m(x)(t - t_0)^{m\beta}, \quad x \in I, \quad 0 \leq m - 1 < \beta \leq m, \quad t_0 \leq t < t_0 + R.$$

If $D_t^{m\beta} u(x, t)$ are continuous on $I \times (t_0, t_0 + R)$, then $g_m(x)$ are expressed as

$$g_m(x) = \frac{D_t^{m\beta} u(x, t_0)}{\Gamma(m\beta + 1)}, \quad m = 0, 1, 2, \dots$$

Here, $D_t^{m\beta} = \frac{\partial^{m\beta}}{\partial t^{m\beta}} = \frac{\partial^\beta}{\partial t^\beta} \cdot \frac{\partial^\beta}{\partial t^\beta} \dots \frac{\partial^\beta}{\partial t^\beta}$, and $R = \min_{c \in I} R_c$ that R_c is radius of convergence of the fractional power series $\sum_{m=0}^{\infty} g_m(c)(t - t_0)^{m\beta}$.

BASIC IDEA OF THE RPSM

In this section, to demonstrate the basic idea of the RPSM, we examine a general nonlinear fractional differential equation by the initial condition of the manner

$$\begin{aligned} D_t^\beta u(x, t) &= N(u) + R(u), \quad 0 < \beta \leq 1, \quad t > 0, \\ u(x, 0) &= g(x), \end{aligned} \tag{3}$$

where D_t^β represents the fractional derivative in the Caputo sense, N is nonlinear differential operator and R is linear differential operator. This method suggests the solution for Eq. (3) as a fractional power series for $t = 0$. Assume the solution takes the following form:

$$u(x, t) = \sum_{m=0}^{\infty} g_m(x) \frac{t^{m\beta}}{\Gamma(m\beta + 1)}, \quad x \in I, \quad 0 < \beta \leq 1, \quad 0 \leq t < R.$$

The $u_l(x, t)$ is also expressed as

$$u_l(x, t) = \sum_{m=0}^l g_m(x) \frac{t^{m\beta}}{\Gamma(m\beta + 1)}, \quad x \in I, \quad 0 < \beta \leq 1, \quad 0 \leq t < R. \tag{4}$$

Then, the 0-th RPS approximate solution of $u(x, t)$ is given as

$$u_0 = g_0(x) = u(x, 0) = g(x).$$

Eq. (4) can be written as

$$u_l(x, t) = g(x) + \sum_{m=1}^l g_m(x) \frac{t^{m\beta}}{\Gamma(m\beta + 1)}, \quad x \in I, \quad 0 < \beta \leq 1, \quad 0 \leq t < R, \quad l = 1, 2, \dots \tag{5}$$

The residual function for Eq. (3) is expressed as

$$Res_u(x, t) = D_t^\beta u(x, t) - N(u) - R(u).$$

Therefore, $Res_{u,l}$ is stated as

$$Res_{u,l}(x, t) = D_t^\beta u_l(x, t) - N(u_l) - R(u_l). \tag{6}$$

Some significant relations of the suggested method are as follows and it can be seen in [38-42].

$$\begin{aligned} Res_u(x, t) &= 0, \\ \lim_{l \rightarrow \infty} Res_{u,l}(x, t) &= Res_u(x, t) \text{ with } t \geq 0 \text{ and } x \in I, \\ D_t^{m\beta} Res_u(x, 0) &= D_t^{m\beta} Res_{u,l}(x, 0) = 0, \quad m = 0, 1, \dots, l. \end{aligned} \quad (7)$$

Substituting the $u_l(x, t)$ in Eq. (6) and calculating the $D_t^{(l-1)\beta}$ of $Res_{u,l}(x, t)$ for $l = 1, 2, \dots$, the suggested method is clearly expressed. Then, applying the relation (7), the following equation

$$D_t^{(l-1)\beta} Res_{u,l}(x, 0) = 0, \quad 0 < \beta \leq 1, \quad 0 \leq t < R, \quad t = 0, \quad l = 1, 2, \dots \quad (8)$$

is solved to obtain the $g_m(x)$ with $m = 1, 2, \dots, l$ in Eq. (5).

APPROXIMATE SOLUTIONS OF THE FRACTIONAL HARRY DYM EQUATION BY RPSM

In this segment of the study, we utilize the RPSM to gain the RPS solutions for Eq. (1) by the initial condition (2).

Let us consider the residual function for Eq. (1) as

$$Res_u(x, t) = D_t^\beta u(x, t) - u^3(x, t) \frac{\partial^3}{\partial x^3} u(x, t).$$

Therefore, $Res_{u,l}(x, t)$ is written as

$$Res_{u,l}(x, t) = D_t^\beta u_l(x, t) - u_l^3(x, t) \frac{\partial^3}{\partial x^3} u_l(x, t). \quad (9)$$

To determine the $g_1(x)$, we write $l = 1$ in Eq. (9) and we have

$$Res_{u,1}(x, t) = D_t^\beta u_1(x, t) - u_1^3(x, t) \frac{\partial^3}{\partial x^3} u_1(x, t).$$

From Eq. (5) for $l = 1$, we get

$$u_1(x, t) = g(x) + g_1(x) \frac{t^\beta}{\Gamma(\beta + 1)}.$$

Hence,

$$Res_{u,1}(x, t) = g_1(x) - \left(g(x) + g_1(x) \frac{t^\beta}{\Gamma(\beta + 1)} \right)^3 \left(g'''(x) + g_1'''(x) \frac{t^\beta}{\Gamma(\beta + 1)} \right).$$

From Eq. (8), we find the $Res_{u,1}(x, 0) = 0$, and therefore

$$g_1(x) = - \frac{1}{\left(4 - \frac{3}{2}x \right)^{\frac{1}{3}}}.$$

Thus, we get

$$u_1(x, t) = \left(4 - \frac{3}{2}x \right)^{\frac{2}{3}} - \frac{1}{\left(4 - \frac{3}{2}x \right)^{\frac{1}{3}}} \frac{t^\beta}{\Gamma(\beta + 1)}.$$

To determine $g_2(x)$, we write $l = 2$ in Eq. (9) and we have

$$Res_{u,2}(x, t) = D_t^\beta u_2(x, t) - u_2^3(x, t) \frac{\partial^3}{\partial x^3} u_2(x, t).$$

From Eq. (5) at $l = 2$, we get

$$u_2(x, t) = g(x) + g_1(x) \frac{t^\beta}{\Gamma(\beta + 1)} + g_2(x) \frac{t^{2\beta}}{\Gamma(2\beta + 1)}.$$

Hence,

$$\begin{aligned} Res_{u,2}(x, t) &= g_1(x) + g_2(x) \frac{t^\beta}{\Gamma(\beta + 1)} \\ &\quad - \left(g(x) + g_1(x) \frac{t^\beta}{\Gamma(\beta + 1)} + g_2(x) \frac{t^{2\beta}}{\Gamma(2\beta + 1)} \right)^3 \left(g'''(x) + g_1'''(x) \frac{t^\beta}{\Gamma(\beta + 1)} \right. \\ &\quad \left. + g_2'''(x) \frac{t^{2\beta}}{\Gamma(2\beta + 1)} \right). \end{aligned}$$

From Eq. (8), we find $D_t^\beta Res_{u,2}(x, 0) = 0$, and therefore

$$g_2(x) = -\frac{1}{2\left(4 - \frac{3}{2}x\right)^{\frac{4}{3}}}.$$

Thus,

$$u_2(x, t) = \left(4 - \frac{3}{2}x\right)^{\frac{2}{3}} - \frac{1}{\left(4 - \frac{3}{2}x\right)^{\frac{1}{3}} \Gamma(\beta + 1)} \frac{t^\beta}{2} - \frac{1}{2\left(4 - \frac{3}{2}x\right)^{\frac{4}{3}} \Gamma(2\beta + 1)} \frac{t^{2\beta}}{4}.$$

To determine $g_3(x)$, we write $l = 3$ in Eq. (9) and we get

$$Res_{u,3}(x, t) = D_t^\beta u_3(x, t) - u_3^3(x, t) \frac{\partial^3}{\partial x^3} u_3(x, t).$$

From Eq. (5) at $l = 3$, we have

$$u_3(x, t) = g(x) + g_1(x) \frac{t^\beta}{\Gamma(\beta + 1)} + g_2(x) \frac{t^{2\beta}}{\Gamma(2\beta + 1)} + g_3(x) \frac{t^{3\beta}}{\Gamma(3\beta + 1)}.$$

Thus,

$$\begin{aligned} Res_{u,3}(x, t) &= g_1(x) + g_2(x) \frac{t^\beta}{\Gamma(\beta + 1)} + g_3(x) \frac{t^{2\beta}}{\Gamma(2\beta + 1)} \\ &\quad - \left(g(x) + g_1(x) \frac{t^\beta}{\Gamma(\beta + 1)} + g_2(x) \frac{t^{2\beta}}{\Gamma(2\beta + 1)} + g_3(x) \frac{t^{3\beta}}{\Gamma(3\beta + 1)} \right)^3 \left(g'''(x) \right. \\ &\quad \left. + g_1'''(x) \frac{t^\beta}{\Gamma(\beta + 1)} + g_2'''(x) \frac{t^{2\beta}}{\Gamma(2\beta + 1)} + g_3'''(x) \frac{t^{3\beta}}{\Gamma(3\beta + 1)} \right). \end{aligned}$$

From Eq. (8), we gain $D_t^{2\beta} Res_{u,3}(x, 0) = 0$, and therefore

$$g_3(x) = -\frac{1}{\left(4 - \frac{3}{2}x\right)^{\frac{7}{3}}}.$$

Hence,

$$u_3(x, t) = \left(4 - \frac{3}{2}x\right)^{\frac{2}{3}} - \frac{1}{\left(4 - \frac{3}{2}x\right)^{\frac{1}{3}} \Gamma(\beta + 1)} \frac{t^\beta}{2} - \frac{1}{2\left(4 - \frac{3}{2}x\right)^{\frac{4}{3}} \Gamma(2\beta + 1)} \frac{t^{2\beta}}{4} - \frac{1}{\left(4 - \frac{3}{2}x\right)^{\frac{7}{3}} \Gamma(3\beta + 1)} \frac{t^{3\beta}}{7}.$$

Using the same operation for $l = 4$, we get

$$g_4(x) = -\frac{7}{2\left(4 - \frac{3}{2}x\right)^{\frac{10}{3}}}.$$

$$u_4(x, t) = \left(4 - \frac{3}{2}x\right)^{\frac{2}{3}} - \frac{1}{\left(4 - \frac{3}{2}x\right)^{\frac{1}{3}} \Gamma(\beta + 1)} t^\beta - \frac{1}{2 \left(4 - \frac{3}{2}x\right)^{\frac{4}{3}} \Gamma(2\beta + 1)} t^{2\beta} - \frac{1}{\left(4 - \frac{3}{2}x\right)^{\frac{7}{3}} \Gamma(3\beta + 1)} t^{3\beta} - \frac{1}{2 \left(4 - \frac{3}{2}x\right)^{\frac{10}{3}} \Gamma(4\beta + 1)} t^{4\beta}.$$

In Table 1, the $u_4(x, t)$ solution is gained for $\beta = 0.25, \beta = 0.50, \beta = 0.75,$ and $\beta = 1$ with the different values of t and x . Besides, the exact solution is compared with the $u_4(x, t)$ solution for $\beta = 1$ in this table. From Table 1, it can be seen that the absolute error gets smaller as the value of t decreases.

Table 1. Comparing the $u_4(x, t)$ solution and the exact solution with the different values of t and x .

x	t	$\beta = 0.25$	$\beta = 0.50$	$\beta = 0.75$	$\beta = 1$		
		$u_4(x, t)$	$u_4(x, t)$	$u_4(x, t)$	$u_4(x, t)$	Exact solution	Absolute error
-10	0.2	6.83852	6.92921	6.99774	7.04522	7.04522	1.35745×10^{-10}
	0.4	6.78388	6.84877	6.91334	6.96966	6.96966	4.39464×10^{-9}
	0.6	6.74698	6.78650	6.83880	6.89370	6.89370	3.37672×10^{-8}
	0.8	6.71828	6.73363	6.76987	6.81731	6.81731	1.44003×10^{-7}
	1	6.69446	6.68676	6.70475	6.7405	6.7405	4.44810×10^{-7}
-5	0.2	4.75697	4.86719	4.94936	5.00586	5.00586	1.20481×10^{-9}
	0.4	4.69016	4.77026	4.84856	4.91607	4.91607	3.93112×10^{-8}
	0.6	4.64474	4.69472	4.75904	4.82544	4.82544	3.04515×10^{-7}
	0.8	4.60926	4.63019	4.67584	4.73396	4.73396	1.30959×10^{-6}
	1	4.57968	4.57266	4.59684	4.64159	4.64159	4.08059×10^{-6}
0	0.2	1.99562	2.18286	2.30909	2.39222	2.39222	1.21435×10^{-7}
	0.4	1.87560	2.02852	2.15777	2.26110	2.26110	4.12452×10^{-6}
	0.6	1.78945	1.90202	2.01889	2.12609	2.12605	3.33870×10^{-5}
	0.8	1.71927	1.78884	1.88536	1.98673	1.98658	1.50729×10^{-4}
	1	1.65869	1.68342	1.75398	1.84251	1.84202	4.95747×10^{-4}
5	0.2	2.75042	2.62131	2.51370	2.43513	2.43513	1.92919×10^{-7}
	0.4	2.82387	2.74355	2.64987	2.56166	2.56167	5.83621×10^{-6}
	0.6	2.87013	2.83355	2.76562	2.68511	2.68515	4.20345×10^{-5}
	0.8	2.90395	2.90641	2.86895	2.80569	2.80586	1.68489×10^{-4}
	1	2.93043	2.96788	2.96320	2.92353	2.92402	4.90352×10^{-4}
10	0.2	5.26849	5.16913	5.09107	5.03561	5.03561	1.40554×10^{-9}
	0.4	5.32758	5.25941	5.18845	5.12435	5.12435	4.41275×10^{-8}
	0.6	5.36687	5.32792	5.27289	5.21232	5.21232	3.28892×10^{-7}
	0.8	5.39709	5.38516	5.34972	5.29956	5.29956	1.36081×10^{-6}
	1	5.42196	5.43522	5.42122	5.38608	5.38609	4.07902×10^{-6}

For $0 \leq t \leq 1$ and $-30 \leq x \leq 0$ at $\beta = 1$, the comparison of the $u_4(x, t)$ and the exact solution is illustrated in Figure 1. When equal parameters are used, it is seen that the $u_4(x, t)$ solution has nearly the same shape as the exact solution in this figure.




Figure 1. The plot of the $u_4(x, t)$ and exact solution.

In Figure 2, the geometrical behavior of the $u_4(x, t)$ with 3D plot for $0 \leq t \leq 5$, $0 \leq x \leq 1$, and the different values of β is illustrated by suggested method. Besides, the same solution with 2D plot for $t = 5$ and $-10 \leq x \leq 10$ is demonstrated in Figure 3. The solution at $\beta = 0.25$ is showed with the blue line, the solution at $\beta = 0.50$ is showed with the orange line, the solution at $\beta = 0.75$ is showed with the green line, and the solution at $\beta = 1$ is showed with the red line in this figure. All plots in figures are illustrated by the aid of Mathematica 11.3.




Figure 2. 3D plot of the $u_4(x, t)$: (a) $u_4(x, t)$ for $\beta = 0.25$, (b) $u_4(x, t)$ for $\beta = 0.50$, (c) $u_4(x, t)$ for $\beta = 0.75$, (d) $u_4(x, t)$ for $\beta = 1$.




Figure 3. 2D plot of the $u_4(x, 5)$ for the different values of β .

For $\beta = 1$, the third order term solution $u_3(x, t)$ of the RPSM, homotopy perturbation Sumudu transform method (HPSTM) [22], Adomian decomposition method (ADM) [22], and exact solution are compared in Table 2. It is observed from this table that the RPSM solution performed a high accuracy agreement with the ADM and HPSM solution. It is also seen that the accuracy increases as the order of the solution increases.

Table 2. Comparison of HPSTM, ADM, RPSM, and exact solution for $\beta = 1$.

x	t	RPSM	HPSTM [22]	ADM [22]	Exact Solution
0	1	1.843946953	1.843946953	1.843946953	1.842015749
0.2	1	1.694117376	1.694117377	1.694117377	1.691538112
0.4	1	1.5337581542	1.537581542	1.537581542	1.534036644
0.6	1	1.373028020	1.373028020	1.373028020	1.367980757
0.8	1	1.198654865	1.198654865	1.198654865	1.91138425
1	1	1.011880652	1.011880649	1.011880649	1.000000000

CONCLUSIONS

In this study, the RPSM was utilized for obtaining the approximate solutions of Eq. (1). These solutions were illustrated by numerically and graphically for the different values of β , t and x . By comparing the approximate solution and the exact solution, the accuracy and efficiency of the suggested method were demonstrated. When equal parameters were selected, it was observed that the approximate solution had almost the same shape as the exact solution. The proposed method was compared numerically with the HPSTM and the ADM by table. . It was seen from this table that the RPSM made a good agreement with this methods. It is seen from the approximate solutions that only a few iterates were used by the proposed method. With these iterates, an infinite series solutions can be found. The accuracy of the RPSM increases as the order of these solutions increases. Besides, this method does not need a lot of time and computer memory. The RPSM indicates strong performance with less computation than other methods in the literature. Moreover, the RPSM does not require transformation, linearization, discretization, or perturbation. Furthermore, the suggested method can be used to get approximate solutions of different kinds of fractional partial differential equations.

REFERENCES

- [1] Vasconcelos, G. L. and Kadanoff, L. P., (1991) Stationary solutions for the Saffman-Taylor problem with surface tension, *Physical Review A*, 44, (10), 1-6.
- [2] Kelleche, A. and Tatar, N.-E., (2017) Control and exponential stabilization for the equation of an axially moving viscoelastic strip, *Mathematical Methods in the Applied Sciences*, 40, (18), 6239-6253.
- [3] Kelleche, A., (2017) Boundary control and stabilization of an axially moving viscoelastic string under a boundary disturbance, *Mathematical Modelling and Analysis*, 22, (6), 763-784.
- [4] Kelleche, A. and Tatar, N.-E., (2017) Existence and stabilization of a Kirchhoff moving string with a distributed delay in the boundary feedback, *Mathematical Modelling of Natural Phenomena*, 12, (6), 106-117.
- [5] Kelleche, A. and Tatar, N.-E., (2021) Adaptive boundary stabilization of a nonlinear axially moving string, *ZAMM - Journal of Applied Mathematics and Mechanics*, Doi: 10.1002/zamm.202000227, 1-14.
- [6] Hereman, W., Banerjee, P. P., and Chatterjee, M. R., (1989) On the nonlocal equations and nonlocal charges associated with the Harry-Dym hierarchy Korteweg-de Vries equation, *Journal of Physics A: Mathematical and Theoretical*, 22, 241-252.
- [7] Haghghatdoost, G. and Bazghandi, M., (2021) Differential invariants of Harry-Dym equation, *The 11th seminar on geometry and topology Yasouj University*, 1-4.
- [8] Mokhtari, R., (2011) Exact Solutions of the Harry-Dym Equation, *Communications in Theoretical Physics*, 55, 204-208.
- [9] Tian, K., Cui, M., and Liu, Q. P., (2022) A note on Bäcklund transformations for the Harry Dym equation, *Partial Differential Equations in Applied Mathematics*, 5, Article ID: 100352, 1-3.
- [10] González-Gaxiola, O., Ruiz de Chávez, J., and Edeki, S. O., (2018) Iterative method for constructing analytical solutions to the Harry-Dym initial value problem, *International Journal of Applied Mathematics*, 31, (4), 627-640.
- [11] Singh, I. and Kumar, S., (2018) Haar wavelet methods for numerical solutions of Harry Dym (HD), BBM Burger's and 2D diffusion equations, *Bulletin of the Brazilian Mathematical Society*, 49, 313-338.
- [12] Soltani, D. and Khorshidi, M. A., (2013) Application of homotopy perturbation and reconstruction of variational iteration methods for Harry Dym equation and compared with exact solution, *International Journal of Multidisciplinary and Current Research*, 166-169.
- [13] Halim, A. A., (2008) Soliton solutions of the (2+1)-dimensional Harry Dym equation via Darboux transformation, *Chaos, Solitons and Fractals*, 36, 646-653.
- [14] Xiao, Y. and Fan, E., (2019) Long time behavior and soliton solution for the Harry Dym equation, *Journal of Mathematical Analysis and Applications*, *Journal of Mathematical Analysis and Applications*, 480, Article ID: 123248, 1-29.
- [15] Ahmad, B. and Nieto, J. J., (2009) Existence of solutions for nonlocal boundary value problems of higher-order nonlinear fractional differential equations, *Hindawi Abstract and Applied Analysis*, Article ID: 494720, 1-9.
- [16] Wang, Y., Liang, S., and Wang, Q., (2018) Existence results for fractional differential equations with integral and multi-point boundary conditions, *Boundary Value Problems*, 4, 1-11.
- [17] Şenol, M. and Ata, A., (2018) Approximate solution of time-fractional KdV equations by residual power series method, *Journal of Balıkesir University Institute of Science and Technology*, 20, (1), 430-439.
- [18] Akram, G., Sadaf, M., Abbas, M., Zainab, I., and Gillani, S. R., (2022), Efficient techniques for traveling wave solutions of time-fractional Zakharov-Kuznetsov equation, *Mathematics and Computers in Simulation*, 193, 607-622.
- [19] Qurashi, M. M. A., Korpınar, Z., Baleanu, D., and Inc, M., (2017) A new iterative algorithm on the time-fractional Fisher equation: Residual power series method, *Advances in Mechanical Engineering*, 9, (9), 1-8.
- [20] Korpınar, Z., (2017) The residual power series method for solving fractional Klein-Gordon equation, *Sakarya University Journal of Science*, 21, (3), 285-293.
- [21] Al-Khaled, K. and Alquran, M., (2014) An approximate solution for a fractional model of generalized Harry Dym equation, *Mathematical Sciences*, 8, 125-130.

- [22] Kumar, D., Singh, J., and Kılıçman, A., (2013) An efficient approach for fractional Harry Dym equation by using Sumudu transform, *Hindawi Abstract and Applied Analysis*, Article ID: 608943, 1-8.
- [23] Alshammari, S., Iqbal, N., and Yar, M., (2022) Analytical Investigation of Nonlinear Fractional Harry Dym and Rosenau-Hyman Equation via a Novel Transform, *Hindawi Journal of Function Spaces*, Article ID: 8736030, 1-12.
- [24] Assabaai, M. A. and Mukherij, O. F., (2020) Exact solutions of the Harry Dym Equation using Lie group method, *University of Aden Journal of Natural and Applied Sciences*, 24, (2), 487-493.
- [25] Costa, F. S., Soares, J. C. A., Plata, A. R. G., and Oliveira, E. C., (2018) On the fractional Harry Dym equation, *Computational and Applied Mathematics*, 37, 2862-2876.
- [26] Yue, C., Liu, G., Li, K., and Dong, H., (2021) Similarity solutions to nonlinear diffusion/Harry Dym fractional equations, *Hindawi Advances in Mathematical Physics*, Article ID: 6670533, 1-6.
- [27] Ghiasi, E. K. and Saleh, R., (2017) A mathematical approach based on the homotopy analysis method: Application to solve the nonlinear Harry-Dym (HD) equation, *Applied Mathematics*, 8, 1546-1562.
- [28] Shunmugarajan, B., (2016) An efficient approach for fractional Harry Dym equation by using homotopy analysis method, *International Journal of Engineering Research & Technology*, 5, (1), 561-566.
- [29] Huang, Q. and Zhdanov, R., (2014) Symmetries and exact solutions of the time fractional Harry-Dym equation with Riemann-Liouville derivative, *Physica A*, 409, 110-118.
- [30] Kumar, S., Tripathi, M. P., Singh, O. P., (2013) A fractional model of Harry Dym equation and its approximate solution, *Ain Shams Engineering Journal*, 4, 111-115.
- [31] Nadeem, M., Li, Z., and Alsayyad, Y., (2022) Analytical approach for the approximate solution of Harry Dym equation with Caputo fractional derivative, *Hindawi Mathematical Problems in Engineering*, Article ID: 4360735, 1-7.
- [32] Rawashdeh, M. S., (2014) A new approach to solve the fractional Harry Dym equation using the FRDTM, *International Journal of Pure and Applied Mathematics*, 95, (4), 553-566.
- [33] Yokuş, A. and Gülbahar, S., (2019) Numerical solutions with linearization techniques of the fractional Harry Dym equation, *Applied Mathematics and Nonlinear Sciences*, 4, (1), 35-42.
- [34] Iyiola, O. S. and Gaba, Y. U., (2016) An analytical approach to time-fractional Harry Dym equation, *Applied Mathematics and Information Sciences*, 10, (2), 409-412.
- [35] Wang, L., Wang, D. Shen, S. and Huang, Q., (2018) Lie point symmetry analysis of the Harry-Dym type equation with Riemann-Liouville fractional derivative, *Acta Mathematicae Applicatae Sinica, English Series*, 34, (3), 469-477.
- [36] Podlubny, I., (1999) Fractional differential equations, New York: Academic Press.
- [37] El-Ajou A., Arqub, O. A., Zhou, Z. A., and Momani, S., (2013) New results on fractional power series: theories and applications, *Entropy*, 15, 5305-5323.
- [38] El-Ajou, A., Arqub, O. A., and Momani, S., (2015) Approximate analytical solution of the nonlinear fractional KdV-Burgers equation: A new iterative algorithm, *Journal of Computational Physics*, 293, 81-95.
- [39] Arqub, A., (2013), Series solution of Fuzzy differential equations under strongly generalized differentiability, *Journal of Advanced Research in Applied Mathematics*, 5,(1), 31-52.
- [40] Arqub, O. A., Abo-Hammour, Z., Al-Badarneh, R., and Momani, S., (2013) A reliable analytical method for solving higher-order initial value problems, *Hindawi Discrete Dynamics in Nature and Society*, Article ID: 673829, 1-12.
- [41] Arqub, O. A., El-Ajou, A., Zhou, Z. A., and Momani, S., (2014) Multiple solutions of nonlinear boundary value problems of fractional order: A new analytic iterative technique, *Entropy*, 16, 471-493.
- [42] Arqub, O. A., El-Ajou, A., Bataineh, A. S., and Hashim, I., (2013) A representation of the exact solution of generalized Lane-Emden equations using a new analytical method, *Hindawi Abstract and Applied Analysis*, Article ID: 378593, 1-10.