

# Sigma Journal of Engineering and Natural Sciences

Web page info: https://sigma.yildiz.edu.tr DOI: 10.14744/sigma.2025.00155



# **Research Article**

# Exploration of Ulam-Hyers stability for a system of fractional integrocoupled differential equations with integral boundary conditions

Brijendra Kumar CHAURASIYA<sup>1</sup>, Avadhesh KUMAR<sup>1,\*</sup>

<sup>1</sup>Department of Mathematics, School of Advanced Sciences, VIT-AP University, Amaravati, 522237, India

#### **ARTICLE INFO**

Article history Received: 16 May 2024 Revised: 03 July 2024 Accepted: 04 October 2024

#### **Keywords:**

Coupled System; Fractional Integro-Differential Equation, Green's Function; Integral Boundary Conditions; Ulam Stability

# **ABSTRACT**

This study examines existence, uniqueness, and Ulam-Hyers stability for solutions of nonlinear coupled fractional integro-differential equations with integral boundary conditions. Fractional systems incorporating memory and hereditary effects serve as effective models for complex processes in science and engineering applications. This study proves solution existence and uniqueness by applying the Banach fixed-point theorem within carefully constructed function spaces, then extends our analysis to investigate Ulam-Hyers stability, a framework that reveals how solutions behave when initial conditions contain small errors. Our stability analysis demonstrates that minor perturbations in starting data translate to bounded solution variations, keeping the system stable within predictable limits, which we verify through a computational example showing how controlled initial changes produce correspondingly controlled solution deviations. These results advance stability theory for coupled fractional integro-differential systems, particularly where memory effects influence system behavior, mathematical models that appear frequently in applications ranging from viscoelastic materials to population dynamics, where past states influence current evolution. By establishing rigorous stability bounds, our work provides a theoretical foundation for implementing these models in real-world scenarios where measurement uncertainties and modeling approximations are inevitable.

Cite this article as: Chaurasiya BK, Kumar A. Exploration of Ulam-Hyers stability for a system of fractional integro-coupled differential equations with integral boundary conditions. Sigma J Eng Nat Sci 2025;43(5):1643–1652.

# INTRODUCTION

Fractional calculus has proven remarkably useful across mathematics and its applications. What makes it fascinating is how we can generalize traditional calculus beyond integer orders, taking derivatives of order 0.5 or 1.7, for instance. This flexibility reveals new mathematical structures that

better model phenomena where history matters, opening doors to more accurate descriptions of complex real-world systems. And it began when L'Hospital and Leibniz wrote a letter to each other in 1695. At present, there are a lot of theories and numerical findings on this topic in the scientific literature. The flow of groundwater in a stream can be described through a fractional differential equation

This paper was recommended for publication in revised form by Editor-in-Chief Ahmet Selim Dalkilic



<sup>\*</sup>Corresponding author.

<sup>\*</sup>E-mail address: soni.iitkgp@gmail.com

that accounts for the non-integer order derivatives. The fractional derivative models the anomalous behavior of flow and transport in porous media. It may also be used in several fields, such as the theory of control, aerodynamics, communication and image filtering, and biology [1-4]. Manuals by Kilbas et al. [5], Miller and Ross [6], and Halfer were used to clarify the fundamental concepts of fractional calculus. Multiple writers have researched the existence and uniqueness of approaches to fractional order differential systems in finite and infinite-dimensional domains. Ahmad et al. discovered results involving non-linear fractional integro-differential equations utilizing integral boundary conditions [7-10].

The results obtained in this research demonstrate practical utility across multiple domains where complex differential systems with integral boundary conditions govern physical phenomena. Hydrogeological researchers have found fractional integro-differential equations particularly effective for modeling subsurface water flow dynamics [10], while chemical engineers apply these systems to understand reaction-diffusion mechanisms [11,12]. When studying population dynamics or circulatory systems, we need mathematical frameworks that can reliably predict how these systems evolve over time, making solution stability absolutely essential for any practical model [13]. Our Ulam-Hyers stability analysis tackles this challenge by establishing clear bounds on how solutions change when initial parameters vary slightly. This becomes especially important in hydrology and biomedical engineering, where professionals rely on model predictions to make decisions that can have serious consequences for public health or environmental protection. The stability criteria we've developed offer mathematical confidence that these models will behave consistently under real-world conditions, which is exactly what practitioners need when applying these tools to solve actual problems in environmental science and engineering.

Integral boundary conditions play a crucial role in hydrogeological modeling as they capture the dynamic interactions between aquifer systems and surrounding environments. These conditions can represent various physical processes, including groundwater recharge rates or discharge flows between aquifers and surface water bodies such as rivers and lakes. Additional details regarding integral boundary conditions can be found in [13-15]. To illustrate the practical significance of integral boundary conditions, consider the following real-world example

$$-\varpi'' = f(\phi)g(\phi, \varpi), \ \varpi(a) = 0, \ \eta\varpi'(b) = \varpi(\beta)$$

where  $\phi \in (a, b)$ ,  $\beta \in (0,1]$  and  $\eta$  are positive constants, and a, b are constants. This represents a thermoregulator model. The problem admits well-defined solutions for the one-dimensional heat equation describing a heated bar equipped with a control system that modulates heat input based on temperature measurements from a sensor positioned at  $\beta$ .

This formulation extends to more general cases as the heat equation, incorporating nonlinear gradient terms and time-varying source components. The heated bar now features a controller positioned at  $\beta$  that regulates heat output according to temperature data collected from multiple sensors distributed along the bar [16-20]. This problem, therefore, may be stated as follows

$$\varpi'' = g(\phi, \varpi, \varpi'), \varpi(a) = 0, \ \varpi'(b) = \int_a^b \varpi(s)dh(s)(\beta).$$

Ulam's type stability is a mathematical notion that describes how solutions to differential equations behave when tiny perturbations are introduced into the beginning conditions or equations themselves. It quantifies how stable or responsive a system is to disturbances. In 1940, Ulam [21] was the first to introduce the Ulam kind of stability. Hyers [22] subsequently refined this concept. Ulam and Hyers investigated stability properties across various integer-order differential equations, leading to what is now termed Hyers-Ulam stability for Cauchy functional equations. Their work produced stronger results concerning polynomials, isometric mappings, and convex functions [23-34]. Numerous researchers have since extended and generalized Hyers' framework for integer-order differential equations. The literature contains extensive investigations into additional stability concepts of Ulam type, documented in [35-44] and related works [45-51].

Motivated by the previously mentioned work related to integro-systems equipped with fractional order derivatives, we aim to extend the findings of [1, 51]. Integro-differential equations are important because they combine the properties of both differential and integral equations, allowing them to model a wide range of complex systems where current states depend on both rates of change and accumulated effects. Despite their significance, few studies have focused on investigating Ulam-type stability for integrodifferential equations involving fractional derivatives. Despite extensive work on fractional differential equations, Ulam's stability for coupled systems with integral boundary conditions remains unexplored. The proposed system is as follows

$$\begin{cases} {}^{c}D^{m}x(t) = k(t,y(t),\chi^{y}(t)), & m \in (1,2], t \in \mathbb{N} \\ {}^{c}D^{n}y(t) = l(t,x(t),\chi^{x}(t)), & n \in (1,2], t \in \mathbb{N} \\ \theta x(0) + \theta x'(0) = \int_{0}^{1} b_{1}(x(s))ds, \theta x(1) + \theta x'(1) = \int_{0}^{1} b_{2}(x(s))ds & \\ \bar{\theta} y(0) + \bar{\theta} y'(0) = \int_{0}^{1} \bar{b}_{1}(y(s))ds, \bar{\theta} y(1) + \bar{\theta} y'(1) = \int_{0}^{1} \bar{b}_{2}(y(s))ds, & \end{cases}$$
(1.1)

where  ${}^cD$  is the Caputo fractional derivative, m, n represents the order of the derivative,  $\aleph = [0,1]$  and  $k, l : [0,1] \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$  for  $\chi^y, \chi^x : [0,1] \times [0,1] \to [0,\infty)$ . And

$$\chi^{y}(t) = \int_0^t \hat{\gamma}_1(t,s)y(s)ds$$
 and  $\chi^{x}(t) = \int_0^t \hat{\gamma}_2(t,s)x(s)ds$ .

Here  $b_1, b_2, \bar{b}_1, \bar{b}_2 \colon \mathbb{R} \to \mathbb{R}$  and  $\theta, \bar{\theta} > 0, \vartheta, \bar{\vartheta} \ge 0$  are real numbers.

We organize this work as follows: Section 2 presents essential definitions, notation, and preliminary results needed for our main analysis. In Section 3, we establish the existence and uniqueness of solutions for system (1.1). Section 4 develops our main stability results, focusing on Ulam-type stability criteria. Finally, Section 5 demonstrates these theoretical findings through a computational example that illustrates the practical implications of our stability bounds.

#### **PRELIMINARIES**

We start with preliminary results using a few helpful concepts, notations, and lemmas, which will be utilized in the subsequent parts.

**Definition 1.**[5] The Caputo derivative of a given function  $y \in ((0,1), \mathbb{R})$  for a fractional order  $m \in \mathbb{R}^+$  is presented as

$$^{c}D^{m}y(t) = \frac{1}{\Gamma(\varsigma - m)} \int_{0}^{t} (t - s)^{\varsigma - m - 1} y^{(\varsigma)}(s) ds, \varsigma = [m] + 1,$$

where [m] denote the integer part of m and  $\Gamma(\cdot)$  is the gamma function.

**Definition 2.** [5] The Riemann- Liouville fractional derivative of a function y(t) of order m is presented as

$$\mathcal{J}^m y(t) = \frac{1}{\Gamma(\varsigma - m)} \left(\frac{d}{dt}\right)^\varsigma \int_0^t \frac{y(s)}{(t - s)^{m - \varsigma + 1}} ds, \varsigma = [m] + 1,$$

provided the integral exists.

**Lemma 1.**[5] For any real number m > 0, the differential equation

$$^{c}D^{m}v(t)=0$$

has a solution which is presented by

$$y(t) = d_0 + d_1 t + d_2 t^2 + \dots + d_{\varsigma - 1} t^{\varsigma - 1},$$
  
$$d_{\iota} \in \mathbb{R}, \iota = 0, 1, 2, \dots, \varsigma - 1,$$

where  $\varsigma = [m] + 1$ .

**Lemma 2.** [5] The solution for any real number m > 0, of the differential equation

$$^{c}D^{m}y(t) = u(t)$$

will be presented as

$$\mathcal{J}^{m}[{}^{c}D^{m}y(t)] = \mathcal{J}^{m}y(t) + d_{0} + d_{1}t + d_{2}t^{2} + \dots + d_{\varsigma-1}t^{\varsigma-1},$$
  
$$d_{\varsigma} \in \mathbb{R}, \iota = 0,1,2,\dots,\varsigma-1,$$

where  $\varsigma = [m] + 1$ .

**Lemma 3.** [9] For any function  $u, \gamma_1, \gamma_2 \in C([0,1], \mathbb{R})$ , the following boundary value problem has a unique solution of

$$\begin{cases} {}^{c}D^{m}y(t) = u(t), t \in [0,1] \\ \theta y(0) + \vartheta y'(0) = \int_{0}^{1} \gamma_{1}(s)ds, \theta y(1) + \vartheta y'(1) = \int_{0}^{1} \gamma_{2}(s)ds \ \ (2.1) \end{cases}$$

is presented as

$$y(t) = \int_0^1 G_m(t,s)u(s)ds + \frac{1}{\theta^2} \left[ (\theta(1-t) + \vartheta) \int_0^1 \gamma_1(s)ds + (\vartheta + \theta t) \int_0^1 \gamma_2(s)ds \right]. \tag{2.2}$$

where  $G_m(t,s)$  is the Green's function presented by

$$G_m(t,s) = \begin{cases} \frac{\theta(t-s)^{m-1} + (\theta-\theta t)(1-s)^{m-1}}{\theta\Gamma(m)} + \frac{\theta(\theta-\theta t)(1-s)^{m-2}}{\theta^2\Gamma(m-1)}, & s \leq t \\ \frac{(\theta-\theta t)(1-s)^{m-1}}{\theta\Gamma(m)} + \frac{\theta(\theta-\theta t)(1-s)^{m-2}}{\theta^2\Gamma(m-1)}, & t \leq s \end{cases} \tag{2.3}$$

**Proof:** To prove lemma (3) for some real constants  $s_1$ ,  $s_2 \in \mathbb{R}$ , we have

$$y(t) = \mathcal{J}^m u(t) - s_1 - s_2 t = \int_0^t \frac{(t - s)^{m - 1}}{\Gamma(m)} u(s) ds - s_1 - s_2 t$$

Given the relation  ${}^cD^m\mathcal{J}^my(t)=y(t)$  and  $\mathcal{J}^m\mathcal{J}^n=\mathcal{J}^{m+n}y(t)$  for  $m,n>0,y\in L(0,1)$ , we obtain

$$y'(t) = \int_0^t \frac{(t-s)^{m-2}}{\Gamma(m-1)} u(s) ds - s_2$$

Applying the boundary conditions for (2.1), we get the following

$$\begin{split} s_1 &= \frac{1}{\theta^2} \bigg[ \vartheta \int_0^1 \gamma_2(s) ds - (\vartheta + \theta) \int_0^1 \gamma_1(s) ds \bigg] - \frac{\vartheta}{\theta \Gamma(m)} \int_0^1 (1-s)^{m-1} u(s) ds \\ &- \frac{\vartheta^2}{\theta^2 \Gamma(m-1)} \int_0^1 (1-s)^{m-2} u(s) ds \\ s_2 &= \frac{1}{\theta} \bigg[ \int_0^1 \gamma_1(s) ds - \int_0^1 \gamma_2(s) ds \bigg] + \frac{1}{\Gamma(m)} \int_0^1 (1-s)^{m-1} u(s) ds \\ &+ \frac{\vartheta}{\theta \Gamma(m-1)} \int_0^1 (1-s)^{m-2} u(s) ds \end{split}$$

The unique solution of (2.1) is therefore follows as

$$\begin{split} y(t) &= \int_0^t \left[ \frac{\theta(t-s)^{m-1} + (\vartheta - \theta t)(1-s)^{m-1}}{\theta \Gamma(m)} + \frac{\vartheta(\vartheta - \theta t)(1-s)^{m-2}}{\theta^2 \Gamma(n-1)} \right] u(s) ds \\ &+ \int_t^t \left[ \frac{(\vartheta - \theta t)(1-s)^{m-1}}{\theta \Gamma(m)} + \frac{\vartheta(\vartheta - \theta t)(1-s)^{m-2}}{\theta^2 \Gamma(n-1)} \right] u(s) ds \\ &+ \frac{1}{\theta^2} \left[ (\vartheta(1-t) + \vartheta) \int_0^1 \gamma_1(s) ds + (\vartheta + \theta t) \int_0^1 \gamma_2(s) ds \right] \\ &= \int_0^1 G_m(t,s) u(s) ds + \frac{1}{\theta^2} \left[ (\vartheta(1-t) + \vartheta) \int_0^1 \gamma_1(s) ds + (\vartheta + \theta t) \int_0^1 \gamma_2(s) ds \right] \end{split}$$

where  $G_m(t,s)$  is given by (2.3). This is the required proof.

**Lemma 4.** The space  $\mathcal{H} = y(t) \mid y \in C(\aleph)$  is a Banach space under the defined norm  $\|y\|_{\mathcal{H}} = \max_{t \in \mathcal{N}} |y(t)|$ . Similarly, the norm on product space is defined as  $\|(y, \bar{y})\|_{\mathcal{H} \times \mathcal{H}} = \|y\|_{\mathcal{H}} + \|\bar{y}\|_{\mathcal{H}}$ . Obviously  $(\mathcal{H} \times \mathcal{H}, \|(.,.)\|_{\mathcal{H} \times \mathcal{H}})$  is a Banach space. Moreover the cone  $\mathcal{U} \subset \mathcal{H} \times \mathcal{H}$  is defined as

$$\mathcal{U} = (y, \bar{y}) \in \mathcal{H} \times \mathcal{H} \mid y(t) \ge 0, \bar{y}(t) \ge 0.$$

Here system (1.1) is transformed into a fixed point problem. Let  $\mathcal{S}: \mathcal{H} \times \mathcal{H} \to \mathcal{H} \times \mathcal{H}$  be the operator defined as

$$\mathcal{S}(x,y)(t) = \begin{pmatrix} \int_{0}^{1} G_{m}(t,s)k(s,y(s),\chi^{y}(s))ds + \frac{1}{\theta^{2}} \Big[ (\theta(1-t)+\theta) \int_{0}^{1} b_{1}(x(s))ds \Big] \\ + (\theta+\theta t) \int_{0}^{1} b_{2}(x(s))ds \\ \int_{0}^{1} G_{n}(t,s)l(s,x(s),\chi^{x}(s))ds + \frac{1}{\theta^{2}} \Big[ (\bar{\theta}(1-t)+\bar{\theta}) \int_{0}^{1} \bar{b}_{1}(y(s))ds \Big] \\ + (\bar{\theta}+\bar{\theta} t) \int_{0}^{1} \bar{b}_{2}(y(s))ds \end{pmatrix}$$
(2.4)

It follows that the fixed point of the operator coincides with the solution of the proposed coupled system (1.1).

To prove the existence and uniqueness of solutions to system (1.1), we require the subsequent conditions

**(B1)** For 
$$t \in \aleph$$
,  $\exists \mu_1, \mu_2, \mu_3 \in C(\aleph, \mathbb{R}^+)$  such that

$$\left| k \left( t, y(t), \chi^{y}(t) \right) \right| \leq \mu_{1}(t) + \mu_{2}(t) |y(t)| + \mu_{3}(t) |\chi^{y}(t)|,$$
$$\forall y(t) \in C(\aleph, \mathbb{R})$$

with

$$\mu_1^* = \sup_{t \in \mathbb{N}} \mu_1(t), \ \mu_2^* = \sup_{t \in \mathbb{N}} \mu_2(t), \ \mu_3^* = \sup_{t \in \mathbb{N}} \mu_3(t).$$

Similarly, For  $t \in \aleph$ ,  $\exists \nu_1, \nu_2, \nu_3 \in C(\aleph, \mathbb{R}^+)$  such that

$$\left| l(t, x(t), \chi^{x}(t)) \right| \le \nu_{1}(v) + \nu_{2}(t)|x(t)| + \nu_{3}(t)|\chi^{x}(t)|,$$

$$\forall x(t) \in C(\aleph, \mathbb{R})$$

with

$$v_1^* = \sup_{t \in \mathbb{N}} v_1(t), \ v_2^* = \sup_{t \in \mathbb{N}} v_2(t), \ v_3^* = \sup_{t \in \mathbb{N}} v_3(t).$$

**(B2)** For  $t \in \aleph$ ,  $\exists L_{h_1}, L_{h_2}$  such that

$$|b_1(x(t))| \le L_{b_1}|x(t)|$$
 and  $|b_2(x(t))| \le L_{b_2}|x(t)|$ ,  
 $\forall x(t) \in C(\aleph, \mathbb{R}).$ 

Similarly, For  $t \in \aleph$ ,  $\exists L_{\bar{b}_1}, L_{\bar{b}_2}$  such that

$$\left|\bar{b}_1(y(t))\right| \le L_{b_1}|y(t)| \text{ and } \left|\bar{b}_2(y(t))\right| \le L_{\bar{b}_2}|y(t)|,$$
$$\forall y(t) \in C(\aleph, \mathbb{R}).$$

**(B3)** For  $t \in \aleph$ ,  $\exists L_{\hat{\omega}_1}, L_{\hat{\omega}_2}$  such that

$$\int_0^1 |\hat{\gamma}_1(t,s)| ds \le L_{\hat{\omega}_1} \text{ and } \int_0^1 |\hat{\gamma}_2(t,s)| ds \le L_{\hat{\omega}_2}.$$

(B4)

$$\begin{split} \mathcal{M}_1 &= \int_0^1 |G_m(t,s)| \mu_1(s) < \infty, \\ \mathcal{N}_1 &= \int_0^1 |G_m(t,s)| \mu_2(s) ds + L_{\hat{\omega}_1} \int_0^1 |G_m(t,s)| \mu_3(s) ds \\ &\quad + \frac{(\theta + \vartheta) \left( L_{b_1} + L_{b_2} \right)}{\theta^2} < \frac{1}{2} \\ \mathcal{M}_2 &= \int_0^1 |G_n(t,s)| \nu_1(s) < \infty, \\ \mathcal{N}_2 &= \int_0^1 |G_n(t,s)| \nu_2(s) ds + L_{\hat{\omega}_2} \int_0^1 |G_n(t,s)| \nu_3(s) ds \\ &\quad + \frac{(\bar{\theta} + \bar{\vartheta}) \left( L_{\bar{b}_1} + L_{\bar{b}_2} \right)}{\bar{\theta}^2} < \frac{1}{2} \end{split}$$

(B5) For all  $\bar{y} \in C(\aleph, \mathbb{R})$  and for each  $t \in \aleph$  there exist positive real constants  $\Re_k$ ,  $\bar{\Re}_k$ , such that

$$\left| k \left( t, y(t), \chi^{y}(t) \right) - k \left( t, \overline{y}(t), \overline{\chi}^{y}(t) \right) \right| \leq \Re_{k} |y - \overline{y}| + \overline{\Re}_{k} |\chi^{y} - \overline{\chi}^{y}|$$

Similarly,  $\forall x, \bar{x} \in C(\aleph, \mathbb{R})$  and for each  $t \in \aleph$  there exist positive constants  $\Re_t$ ,  $\bar{\Re}_b$ , such that

$$\left|l(t,x(t),\chi^{x}(v))-k(t,\bar{x}(t),\bar{\chi}^{x}(t))\right| \leq \Re_{t}|x-\bar{x}|+\bar{\Re}_{t}|\chi^{x}-\bar{\chi}^{x}|$$

**(B6)** For all  $\bar{x} \in C(\aleph, \mathbb{R})$  and for each  $t \in \aleph$  there exist positive real constants  $\mathfrak{R}_{b_1}$ ,  $\mathfrak{R}_{b_2}$ , such that

$$|b_1(x) - b_1(\bar{x})| \le \Re_{b_1}|x - \bar{x}| \text{ and } |b_2(x) - b_2(\bar{x})| \le \Re_{b_2}|x - \bar{x}|$$

Similarly, For all  $y, \bar{y} \in C(\aleph, \mathbb{R})$  and for each  $t \in \aleph$  there exist positive real constants  $\Re_{\bar{b}_1}$ ,  $\Re_{\bar{b}_2}$ , such that

$$|\bar{b}_1(y) - \bar{b}_1(\bar{y})| \le \Re_{\bar{b}_1} |y - \bar{y}| \text{ and } |\bar{b}_2(y) - \bar{b}_2(\bar{y})| \le \Re_{\bar{b}_2} |y - \bar{y}|$$

(B7) Let (i). 
$$\sigma_1 = \mathcal{K}_{(n\bar{\theta}\bar{\vartheta})} \Re_l + \Re_{\theta\vartheta}$$
, where  $\Re_{\theta\vartheta} = \frac{(\theta+\vartheta)(\Re_{b_1}+\Re_{b_2})}{\theta^2}$ 

and

$$\begin{split} \mathcal{K}_{(n\bar{\theta}\bar{\theta})} &= \max_{t \in [0,1]} \left| \int_0^1 G_n(t,s) ds \right| \\ &= \left| \int_0^t \left[ \frac{\bar{\theta}(t-s)^{(n-1)} + (\bar{\theta}-\bar{\theta})(1-s)^{n-1}}{\theta \Gamma(n)} + \frac{\bar{\theta}(\bar{\theta}-\bar{\theta}t)(1-s)^{n-2}}{\theta^2 \Gamma(n-1)} \right] ds \right| \\ &+ \left| \int_t^1 \left[ \frac{(\bar{\theta}-\bar{\theta}t)(1-s)^{(n-1)}}{\theta \Gamma(n)} + \frac{\bar{\theta}(\bar{\theta}-\bar{\theta}t)(1-s)^{n-2}}{\theta^2 \Gamma(n)} \right] ds \right| \\ &= \frac{1}{\Gamma(n+1)} + \frac{2(\bar{\theta}+\bar{\theta})}{\Gamma(n+1)} + \frac{2(\theta^2+\bar{\theta}\bar{\theta})}{\theta^2 \Gamma(n)} \end{split}$$

(ii). 
$$\sigma_2 = \mathcal{K}_{(m\theta\vartheta)} \Re_k + \Re_{\bar{\theta}\bar{\vartheta}}$$
, where  $\Re_{\bar{\theta}\bar{\vartheta}} = \frac{(\bar{\theta} + \bar{\vartheta})(\Re_{\bar{b}_1} + \Re_{\bar{b}_2})}{\theta^2}$ 

and

$$\mathcal{K}_{(m\theta\vartheta)} = \max_{t \in [0,1]} \left| \int_0^1 G_m(t,s) ds \right|$$
$$= \frac{1}{\Gamma(m+1)} + \frac{2(\theta+\vartheta)}{\Gamma(m+1)} + \frac{2(\vartheta^2 + \theta\vartheta)}{\theta^2 \Gamma(m)}$$

# **EXISTENCE AND UNIQUENESS ANALYSIS**

**Theorem 3.1.** If all of the prerequisites (*B*1) - (*B*7) are satisfied and  $\sigma = \max(\sigma_1, \sigma_2) < 1$ , so the suggested fractional order paired framework (1.1) has a unique solution.

Proof: For a positive number

$$\rho = \max \Big(\frac{2\mathcal{M}_1}{1-2\mathcal{N}_1}, \frac{2\mathcal{M}_2}{1-2\mathcal{N}_2}\Big),$$

we define a set

$$\mathcal{V} = \{(x, y) \in \mathcal{H} \times \mathcal{H} : \| (x, y) \|_{\mathcal{H} \times \mathcal{H}} \leq \rho \}.$$

First, to establish that S maps V into itself, we have

$$\begin{split} |S_m(n,m)(t)| &\leq \int_0^1 |G_m(t,s)| |k(s,y(s),\chi^y(s)) ds| \\ &+ \frac{1}{\alpha^2} \bigg[ |\{\theta(1-t)+\vartheta)| \int_0^1 |b_1(x(s))| ds + |(\vartheta+\theta t)| \int_0^1 |b_2(x(s))| ds \bigg] \\ &\leq \int_0^1 |G_m(t,s)| \mu_1(s) ds + \int_0^1 |G_m(t,s)| |\mu_2(s)| y(s)| ] ds + \int_0^1 |G_m(t,s)| [\mu_3(s)|\chi^y(s)|] ds \\ &+ \frac{1}{\theta^2} \bigg[ (\theta+\vartheta) \int_0^1 |b_1(x(s))| ds + (\theta+\vartheta) \int_0^1 |b_2(x(s))| ds \bigg] \\ &\leq \int_0^1 |G_m(t,s)| \mu_1(s) ds + \rho \bigg[ \int_0^1 |G_m(t,s)| \mu_2(s) ds + L_{\omega_1} \int_0^1 |G_m(t,s)| \mu_3(s) ds \,. \end{split}$$

Upon applying the maximization operation to both sides of the aforementioned inequality over the set  $\mathcal{T}$ , we derive the following expression:

$$\|S_m(n,m)\|_{\mathcal{H}} \leq \frac{\rho}{2}.$$

Similarly,

$$\left\| \mathcal{S}_n(m,n) \right\|_{\mathcal{H}} \le \frac{\rho}{2}.$$

Hence, we can conclude that,

$$\parallel \mathcal{S}(m,n) \parallel_{\mathcal{H} \times \mathcal{H}} \leq \rho. \tag{3.2}$$

The aforementioned inequality serves as confirmation that the mapping S effectively maps the set V into itself. Subsequently, to demonstrate that S qualifies as the contraction operator under the condition where  $t \in \aleph$ , we proceed with the following analysis.

$$\begin{split} \left| \mathcal{S}_m(n,m)(t) - \mathcal{S}_m(\bar{y},\bar{x})(t) \right| &\leq \int_0^1 \left| G_m(t,s) \right| \left| k \left( s, y(s), \chi^y(s) \right) - k \left( s, \bar{y}(s), \bar{\chi}^y(s) \right) \right| ds \\ &+ \frac{1}{\theta^2} \left[ \left| (\theta(1-t) + \theta) \right| \int_0^1 \left| b_1 \left( x(s) \right) - b_1 \left( \bar{x}(s) \right) \right| ds \\ &\leq \mathcal{K}_{(m\theta\theta)} \Re_k |y(t) - \bar{y}(t)| \\ &+ \frac{(\theta + \theta) \left( \Re_{b_1} + \Re_{b_2} \right)}{\theta^2} |x(t) - \bar{x}(t)|. \end{split}$$

When we operate the maximum on both sides of the above-mentioned inequality over  $\aleph$ , we obtain

$$\left\| \mathcal{S}_m(y,x) - \mathcal{S}_m(\bar{y},\bar{x}) \right\|_{\mathcal{H}} \leq \mathcal{K}_{(m\theta\theta)} \Re_k \parallel y - \bar{y} \parallel_{\mathcal{H}} + \frac{(\theta + \vartheta) \left(\Re_{b_1} + \Re_{b_2}\right)}{\theta^2} \parallel x - \bar{x} \parallel_{\mathcal{H}}. \eqno(3.4)$$

Similarly, the following can be obtained

$$\|\mathcal{S}_n(x,y) - \mathcal{S}_n(\bar{x},\bar{y})\|_{\mathcal{H}} \leq \mathcal{H}_{(n\bar{\theta}\bar{\theta})} \mathfrak{R}_{l} \parallel x - \bar{x} \parallel_{\mathcal{H}} + \frac{(\bar{\theta} + \bar{\vartheta}) (\mathfrak{R}_{\bar{h}_1} + \mathfrak{R}_{\bar{h}_2})}{\bar{\theta}^2} \parallel y - \bar{y} \parallel_{\mathcal{H}} \cdot \left(3.5\right)$$

From the above two equations, we obtain

$$\parallel \mathcal{S}(x,y) - \mathcal{S}(\bar{x},\bar{y}) \parallel_{\mathcal{H} \times \mathcal{H}} \leq \sigma \parallel (x,y) - (\bar{x},\bar{y}) \parallel_{\mathcal{H} \times \mathcal{H}}. \quad (3.6)$$

Henceforth, it is evident that the operator  $\mathcal{S}$  exhibits strict contraction properties. Employing Banach's fixed-point method, specifically designed for establishing a unique fixed point, becomes instrumental in determining the sole solution to the proposed coupled system (1.1). Consequently, the fulfillment of our primary objective is achieved.

#### **ULAM'S STABILITY ANALYSIS**

In this section, we endeavor to investigate the Ulamtype stability characteristics inherent in the envisaged coupled system denoted as (1.1).

For some real number  $\varepsilon = \max(\varepsilon_m, \varepsilon_n) > 0$ . We assume the inequality given as follows

$$\begin{cases} \left| {^c}D^m x(t) - k(t, y(t), \chi^y(t)) \right| \le \varepsilon_m, t \in \aleph \\ \left| {^c}D^n y(t) - l(t, x(t), \chi^x(t)) \right| \le \varepsilon_n, t \in \aleph. \end{cases}$$
(4.1)

The following definitions are inspired by Rus [19].

**Definition 3:** The Ulam-Hyers stability of the coupled system denoted by (1.1) is established under the condition that there exists a positive constant  $\mathcal{K}_{(mn\theta\nu\bar{\theta}\bar{\nu})}$  such that, for any solution  $(x, y) \in \mathcal{H} \times \mathcal{H}$  of the inequality (4.1), there exists a unique solution  $(\zeta, \eta) \in \mathcal{H} \times \mathcal{H}$ . This unique solution satisfies the inequality:

$$|(x,y)(t) - (\zeta,\eta)(t)| \le \mathcal{K}_{(mn\theta\vartheta\bar{\vartheta}\bar{\vartheta})}\varepsilon, \quad t \in \aleph$$

**Definition 4:** The generalized Ulam's-Hyers stability of the coupled system denoted by (1.1) is posited under the condition that there exists a function  $\varpi(\varepsilon) \in \mathcal{C}(\mathbb{R}^+, \mathbb{R}^+)$  with  $\varpi(0) = 0$ . This function is required to satisfy the condition that, for any solution  $(x, y) \in \mathcal{H} \times \mathcal{H}$  of the inequality (4.1), there exists a unique solution  $(\zeta, \eta) \in \mathcal{H} \times \mathcal{H}$  for the system. Additionally, the unique solution must adhere to the inequality:

$$|(x,y)(t) - (\zeta,\eta)(t)| \le \varpi(\varepsilon), \quad t \in \aleph$$

**Remark 1.**Consider the pair  $(x, y) \in \mathcal{H} \times \mathcal{H}$  to be a solution of inequality (4.1) if there exist functions  $\phi$  and  $\psi$  belonging to the space  $C(\aleph, \mathbb{R})$ , contingent upon the variables x and y respectively, such that

(R1). 
$$|\phi(t)| \le \varepsilon_m$$
 and  $|\psi(t)| \le \varepsilon_n$ ,  $t \in \aleph$ ;

(R2). and

$$\begin{cases} {}^{c}D^{m}x(t) = k(t, y(t), \chi^{y}(t)) + \phi(t), t \in \aleph \\ {}^{c}D^{n}y(t) = l(t, x(t), \chi^{x}(t)) + \psi(t), t \in \aleph. \end{cases}$$
(4.2)

**Lemma 5.** Let  $(x, y) \in \mathcal{H} \times \mathcal{H}$  be deemed as a solution to inequality (4.1). Subsequently, the ensuing inequality is guaranteed to be satisfied:

$$\left\{ \left| \begin{array}{l} x(t) - \int_0^1 G_m(t,s) k \big( s,y(s),\chi^y(s) \big) ds - \frac{1}{\theta^2} \bigg[ \big( \theta(1-t) + \vartheta \big) \int_0^1 b_1 \big( x(s) \big) ds \\ \\ + \big( \vartheta + \theta t \big) \int_0^1 b_2 \big( x(s) \big) ds \bigg] \right| \leq \mathcal{K}_{m\theta\vartheta} \varepsilon_m, t \in \mathbb{R} \\ \left| \begin{array}{l} y(t) - \int_0^1 G_n(t,s) l \big( s,x(s),\chi^x(s) \big) ds - \frac{1}{\theta^2} \bigg[ \big( \bar{\theta}(1-t) + \bar{\vartheta} \big) \int_0^1 \bar{b}_1 \big( x(s) \big) ds \\ \\ + \big( \bar{\vartheta} + \bar{\theta} t \big) \int_0^1 \bar{b}_2 \big( x(s) \big) ds \bigg] \right| \leq \mathcal{K}_{n\bar{\vartheta}\bar{\vartheta}} \varepsilon_n, t \in \mathbb{R}. \end{array}$$

Proof. By Remark 1 (R2), we have

$$\begin{cases} {}^{c}D^{m}x(t) = k(t, y(t), \chi^{y}(t)) + \phi(t), & t \in \aleph \\ {}^{c}D^{n}y(t) = l(t, x(t), \chi^{x}(t)) + \psi(t), & t \in \aleph \\ \theta x(0) + \vartheta x'(0) = \int_{0}^{1} b_{1}(x(s))ds, \theta x(1) + \vartheta x'(1) = \int_{0}^{1} b_{2}(x(s))ds \\ \bar{\theta} y(0) + \bar{\vartheta} y'(0) = \int_{0}^{1} \bar{b}_{1}(y(s))ds, \bar{\theta} y(1) + \bar{\vartheta} y'(1) = \int_{0}^{1} \bar{b}_{2}(y(s))ds. \end{cases} \tag{4.3}$$

By applying Lemma 3, the solution of (4.3) is given as follows:

$$\begin{cases} x(t) = \int_0^1 G_m(t,s)k\big(s,y(s),\chi^y(s)\big)ds + \int_0^1 G_m(t,s)\phi(s)ds + \frac{1}{\theta^2} \big[(\theta(1-t)+\vartheta) \\ \int_0^1 b_1\big(x(s)\big)ds + (\vartheta+\theta t)\int_0^1 b_2\big(x(s)\big) \bigg]ds, t \in \mathbb{N} \\ y(t) = \int_0^1 G_n(t,s)l\big(s,x(s),\chi^x(s)\big)ds + \int_0^1 G_n(t,s)\psi(s)ds + \frac{1}{\theta^2} \big[(\bar{\theta}(1-t)+\bar{\theta}) \\ \int_0^1 \bar{b}_1\big(x(s)\big)ds + (\bar{\theta}+\bar{\theta}t)\int_0^1 \bar{b}_2\big(x(s)\big) \bigg]ds, t \in \mathbb{N}. \end{cases} \tag{4.4}$$

Considering the first equation of system (4.4), we have

$$\begin{split} \mid x(t) - \int_0^1 G_m(t,s) k\big(s,y(s),\chi^y(s)\big) ds - \frac{1}{\theta^2} \bigg[ \big(\theta(1-t) + \theta\big) \int_0^1 b_1\big(x(s)\big) ds \\ + \big(\theta + \theta t\big) \int_0^1 b_2\big(x(s)\big) \bigg] ds \mid \leq \left| \int_0^1 G_\theta(t,s) \phi(s) ds \right| \\ & \leq \int_0^1 |G_\theta(t,s) \phi(s)| ds. \end{split}$$

By Remark 1 (R1) and using the condition of (B6), we get

$$\begin{vmatrix} x(t) - \int_0^1 G_m(t,s)k(s,y(s),\chi^y(s))ds - \frac{1}{\theta^2} \Big[ (\theta(1-t) + \theta) \int_0^1 b_1(x(s))ds \\ + (\theta + \theta t) \int_0^1 b_2(x(s)) \Big] ds | \leq \mathcal{K}_{(m\theta\theta)} \varepsilon_m.$$
 (4.5)

By applying the same process for the second equation of system (4.4), we get

$$\left| y(t) - \int_0^1 G_n(t,s)l(s,x(s),\chi^*(s))ds - \frac{1}{\bar{\theta}^2} \left[ (\bar{\theta}(1-t) + \bar{\theta}) \int_0^1 \bar{b}_1(x(s))ds + (\bar{\theta} + \bar{\theta}t) \int_0^1 \bar{b}_2(x(s)) \right] ds \right| \le \mathcal{K}_{(n\bar{\theta}\bar{\theta})} \varepsilon_n.$$

$$(4.6)$$

**Theorem 2.** Assuming that conditions (B5) to (B7) are satisfied, the fractional order coupled system denoted by equation (1.1) can be deemed Ulam's-Hyers stable. This

assertion extends to the broader context of being generalized Ulam 's-Hyers stable.

$$(1 - \Re_{\theta\theta})(1 - \Re_{\bar{\theta},\bar{\theta}}) - \Re_{m\theta\theta}\Re_{n\bar{\theta},\bar{\theta}} \neq 0$$

**Proof:** Let  $(x, y) \in \mathcal{H} \times \mathcal{H}$  be the solution of the system (12) and  $(\zeta, \eta) \in \mathcal{H} \times \mathcal{H}$  be the unique solution to the following considered system:

$$\begin{cases} {}^cD^m\zeta(t)=k\big(t,\eta(t),\chi^\eta(t)\big),t\in\aleph\\ {}^cD^n\eta(t)=l\big(t,\zeta(t),\chi^\zeta(t)\big),t\in\aleph\\ \theta\zeta(0)+\vartheta\zeta'(0)=\int_0^1b_1\big(\zeta(s)\big)ds,\theta\zeta(1)+\vartheta\zeta'(1)=\int_0^1b_2\big(\zeta(s)\big)ds\\ \bar\theta\eta(0)+\bar\vartheta\eta'(0)=\int_0^1\overline{b_1}\big(\eta(s)\big)ds,\bar\theta\eta(1)+\bar\vartheta\eta'(1)=\int_0^1\overline{b_2}\big(\eta(s)\big)ds. \end{cases}$$

Using Lemma 3, the solution of the above system is

$$\begin{cases} \zeta(t) = \int_0^1 G_m(t,s) k\big(s,\eta(s),\chi^\eta(s)\big) ds + \frac{1}{\theta^2} \bigg[ \big(\theta(1-t)+\vartheta\big) \int_0^1 b_1\big(\zeta(s)\big) ds \\ \eta(t) = \int_0^1 G_n(t,s) l\left(s,\zeta(s),\chi^\zeta(s)\right) ds + \frac{1}{\bar{\theta}^2} \bigg[ \big(\bar{\theta}(1-t)+\bar{\vartheta}\big) \int_0^1 \bar{b}_1\big(\eta(s)\big) ds \\ + \big(\bar{\vartheta}+\bar{\theta}t\big) \int_0^1 \bar{b}_2\big(\eta(s)\big) \bigg] ds, \quad t \in \aleph. \end{cases}$$

We have

$$\begin{aligned} |x(t) - \zeta(t)| &= \left| x(t) - \int_0^1 \left| G_m(t, s) k \left( s, \eta(s), \chi^{\eta}(s) \right) ds - \frac{1}{\theta^2} \right[ (\theta(1 - t) + \theta) \int_0^1 b_1(\zeta(s)) ds \right. \\ &\quad + (\theta + \theta t) \int_0^1 b_2(\zeta(s)) ds \right] | \\ &\leq \left| x(t) - \int_0^1 \left| G_m(t, s) k \left( s, y(s), \chi^{y}(s) \right) ds - \frac{1}{\theta^2} \left[ (\theta(1 - t) + \theta) \int_0^1 b_1(x(s)) ds \right. \\ &\quad + (\theta + \theta t) \int_0^1 b_2(x(s)) ds \right] | \\ &\quad + \left| \int_0^1 \left| G_m(t, s) k \left( s, y(s), \chi^{y}(s) \right) ds - \int_0^1 \left| G_m(t, s) k \left( s, \eta(s), \chi^{\eta}(s) \right) ds \right| \right. \\ &\quad + \left| \frac{1}{\theta^2} \left[ (\theta(1 - t) + \theta) \int_0^1 b_1(x(s)) ds + (\theta + \theta t) \int_0^1 b_2(x(s)) ds \right] \\ &\quad - \frac{1}{\theta^2} \left[ (\theta(1 - t) + \theta) \int_0^1 b_1(\zeta(s)) ds + (\theta + \theta t) \int_0^1 b_2(\zeta(s)) ds \right] \end{aligned}$$

where  $\mathfrak{R}_{m\theta\vartheta} = \mathcal{K}_{m\theta\vartheta} \mathfrak{R}_{k}$ . Hence, we get

$$(1 - K_{\theta \vartheta}) \parallel x - \zeta \parallel_{\mathcal{H}} \le \mathcal{K}_{(m\theta \vartheta)} \varepsilon_m + \Re_{m\theta \vartheta} \parallel y - \eta \parallel_{\mathcal{H}} (4.9)$$

Similarly, we have

$$(1 - \Re_{\bar{\theta}\bar{\vartheta}}) \parallel y - \eta \parallel_{\mathcal{H}} \leq \mathcal{K}_{(n\bar{\theta}\bar{\vartheta}}\varepsilon_n + \Re_{n\bar{\theta}\bar{\vartheta}} \parallel x - \zeta \parallel_{\mathcal{H}}, (4.10)$$

where  $\mathfrak{R}_{n\bar{\theta}\bar{\vartheta}} = \mathcal{K}_{n\bar{\theta}\bar{\vartheta}}\mathfrak{R}_{l}$ . From (4.9) and (4.10), it can be written as

$$\begin{cases} (1 - \mathfrak{R}_{\theta\theta}) \parallel x - \zeta \parallel_{\mathcal{H}} - \mathfrak{R}_{m\theta\vartheta} \parallel y - \eta \parallel_{\mathcal{H}} \leq \mathcal{K}_{(m\theta\vartheta)} \varepsilon_m \\ (1 - \mathfrak{R}_{\bar{\theta}\bar{\vartheta}}) \parallel y - \eta \parallel_{\mathcal{H}} - \mathfrak{R}_{n\bar{\theta}\bar{\vartheta}} \parallel x - \zeta \parallel_{\mathcal{H}} \leq \mathcal{K}_{(n\bar{\theta}\bar{\vartheta})} \varepsilon_n . \end{cases}$$

The matrix representation of the above equation is as follows

$$\begin{pmatrix} (1-\mathfrak{R}_{\theta\vartheta}) & -\mathfrak{R}_{m\theta\vartheta} \\ -\mathfrak{R}_{n\bar{\theta}\bar{\vartheta}} & (1-\mathfrak{R}_{\bar{\vartheta}\bar{\vartheta}}) \end{pmatrix} \begin{pmatrix} \parallel x-\zeta \parallel_{\mathcal{H}} \\ \parallel y-\eta \parallel_{\mathcal{H}} \end{pmatrix} \leq \begin{pmatrix} \mathcal{K}_{(m\theta\vartheta)}\varepsilon_m \\ \mathcal{K}_{(n\bar{\theta}\bar{\vartheta})}\varepsilon_n \end{pmatrix}.$$

Simplifying the above inequality yields

$$\begin{pmatrix} \parallel x - \zeta \parallel_{\mathcal{H}} \\ \parallel y - \eta \parallel_{\mathcal{H}} \end{pmatrix} \leq \begin{pmatrix} \frac{(1 - \Re_{\bar{\theta}\bar{\vartheta}})}{\Delta} & \frac{\Re_{m\theta\vartheta}}{\Delta} \\ \frac{\Re_{n\bar{\theta}\bar{\vartheta}}}{\Delta} & \frac{(1 - \Re_{\theta\vartheta})}{\Delta} \end{pmatrix} \begin{pmatrix} \mathcal{K}_{(m\theta\vartheta)}\varepsilon_m \\ \mathcal{K}_{(n\bar{\theta}\bar{\vartheta})}\varepsilon_n \end{pmatrix}.$$

where  $\Delta = (1 - \Re_{\theta\theta})(1 - \Re_{\bar{\theta}\bar{\theta}}) - \Re_{m\theta\theta}\Re_{n\bar{\theta}\bar{\theta}} \neq 0$ . Further simplification gives

$$\parallel x - \zeta \parallel_{\mathcal{H}} \leq \frac{(1 - \mathfrak{R}_{\bar{\theta}\bar{\vartheta}})\mathcal{K}_{(m\theta\vartheta)}\varepsilon_{m}}{\Delta} + \frac{\mathfrak{R}_{m\theta\vartheta}\mathcal{K}_{(n\bar{\theta}\bar{\vartheta})}\varepsilon_{n}}{\Delta} \tag{4.12}$$

$$\parallel y - \eta \parallel_{\mathcal{H}} \leq \frac{\Re_{n\bar{\vartheta}\bar{\vartheta}}\mathcal{K}_{(m\theta\vartheta)}\varepsilon_{m}}{\Delta} + \frac{(1 - \Re_{\theta\vartheta})\mathcal{K}_{(n\bar{\theta}\bar{\vartheta}\bar{\vartheta})}\varepsilon_{n}}{\Delta}, \ (4.13)$$

From equations (4.12) and (4.13), we have

$$\begin{split} \parallel x - \zeta \parallel_{\mathcal{H}} + \parallel y - \eta \parallel_{\mathcal{H}} \leq & \frac{(1 - \Re_{\bar{\theta}\bar{\vartheta}})\mathcal{K}_{(m\theta\vartheta)}\varepsilon_{m}}{\Delta} + \frac{\Re_{m\theta\vartheta}\mathcal{K}_{(n\bar{\theta}\bar{\vartheta})}\varepsilon_{n}}{\Delta} \\ & + \frac{\Re_{n\bar{\theta}\vartheta}\mathcal{R}_{(m\theta\vartheta)}\varepsilon_{m}}{\Delta} + \frac{(1 - \Re_{\theta\vartheta})\mathcal{K}_{(n\bar{\vartheta}\bar{\vartheta}}\bar{\varepsilon}\varepsilon_{n}}{\Delta} \end{split}$$

Therefore, we have

$$\parallel (x,y) - (\zeta,\eta) \parallel_{\mathcal{H} \times \mathcal{H}} \leq \mathcal{K}_{(mn\theta\vartheta\bar{\theta}\bar{v})}\varepsilon, \tag{4.14}$$

where  $\varepsilon = \max(\varepsilon_m, \varepsilon_n)$  and

$$\begin{split} \mathcal{K}_{(mn\theta\vartheta\bar{\theta}\bar{\vartheta})} &= \frac{\left(1 - \Re_{\bar{\theta}\bar{\vartheta}}\mathcal{K}_{(m\theta\vartheta)} + \frac{\Re_{m\theta\vartheta}\mathcal{K}_{(n\bar{\theta}\bar{\vartheta})}}{\Delta} \right. \\ &+ \frac{\Re_{n\bar{\vartheta}}\mathcal{R}_{(m\theta\vartheta)}}{\Delta} + \frac{(1 - \Re_{\theta\vartheta})\mathcal{K}_{(n\bar{\theta}\bar{\vartheta})}}{\Delta} \end{split}$$

Therefore, based on the preceding inequality, it can be deduced that the coupled system denoted as (1.1) exhibits Ulam 's-Hyers stability. Additionally, the aforementioned inequality can be expressed in the form:

$$\|(x,y)-(\zeta,\eta)\|_{\mathcal{H}\times\mathcal{H}}\leq \varpi(\varepsilon)$$
, where  $\varpi(0)=0$ .

Through this inequality, it becomes evident that the proposed coupled system (1.1) adheres to the principles of generalized Ulam 's-Hyers stability.

# **APPLICATION**

**Example.** Here, we validate our findings related to existence, uniqueness, and stability by presenting a concrete example. Let the following fractional order connected coupled system as follows:

$$\begin{cases} {}^cD_{-}^{\frac{1}{2}}x(t) = \frac{1}{(t+)^2} \frac{|y(t)|}{1+|y(x)|} + \int_0^t \frac{e^{-(s-t)}}{49} y(s) ds, m \in (0,1], t \in \mathbb{N} \\ {}^cD_{-}^{\frac{1}{2}}y(t) = \frac{1}{100} [t\cos x(t) + x(v)\sin t] + \int_0^t \frac{7}{s^2 + 7^2} x(s) ds, n \in (0,1], t \in \mathbb{N} \\ x(0) + x'(0) = \int_0^1 \frac{|x(s)|}{11 + |x(s)|} ds, x(1) + x'(1) = \int_0^1 \frac{|x(s)|}{13 + |x(s)|} ds \\ y(0) + y'(0) = \int_0^1 \frac{1}{22} [\cos y(s) + \sin y(s)] ds \\ y(1) + y'(1) = \int_0^1 \frac{1}{26} [\cos y(s) + \sin y(s)] ds. \end{cases}$$

By comparing coupled system (5.1) with system (1.1), the following values are derived:

$$m = n = \frac{1}{2}, \theta = \vartheta = \bar{\theta} = \bar{\vartheta} = 1, \Re_{b_1} = \Re_{\bar{b}_1} = \frac{1}{11}$$
  
and  $\Re_{b_2} = \Re_{\bar{b}_2} = \frac{1}{13}$ .

Also,

$$k(t,y(t),\chi^{y}(t)) = \frac{1}{(t+7)^{2}} \frac{|y(t)|}{1+|y(t)|} + \int_{0}^{t} \frac{e^{-(s-t)}}{49} y(s) ds$$

and

$$l(t, x(t), \chi^{x}(t)) = \frac{1}{100} [t \cos x(t) + x(t) \sin t] + \int_{0}^{t} \frac{7}{s^{2} + 7^{2}} x(s) ds,$$

whereas  $|k(t, y, \chi^y) - k(t, \bar{y}, \bar{\chi}^y)| \le \left(\frac{1}{49}\right) ||y - \bar{y}|| + ||\chi^y - \bar{\chi}^y||$  and  $|l(t, x, \chi^x) - l(t, \bar{x}, \bar{\chi}^x)| \le \frac{1}{50} ||x - \bar{x}|| + ||\chi^x - \bar{\chi}^x||$ .

Therefore, (B4) is satisfied with  $\Re_k$ ,  $\bar{\Re}_k = \frac{1}{49}$  and  $\Re_l$ ,  $\bar{\Re}_l = \frac{1}{50}$ . Further, we have

$$\begin{split} &\sigma_1\!=\!\left(\frac{1}{\Gamma(n+1)}\!+\!\frac{2(\bar{\theta}+\bar{\vartheta})}{\Gamma(n+1)}\!+\!\frac{2(\bar{\theta}^2+\bar{\theta}\bar{\vartheta})}{\bar{\theta}^2\Gamma(n)}\right)\Re_l+\!\frac{(\theta+\vartheta)\!\left(\Re_{b_1}\!+\!\Re_{b_2}\right)}{\theta^2}\\ &=\left(\frac{1}{\Gamma(n+1)}\!+\!\frac{4}{\Gamma(n+1)}\!+\!\frac{4}{\Gamma(n)}\right)\!\frac{1}{50}\!+\!2\left(\frac{1}{11}\!+\!\frac{1}{13}\right)\\ &=\left(\frac{5}{\Gamma(n+1)}\!+\!\frac{4}{\Gamma(n)}\right)\!\frac{1}{50}\!+\!\left(\frac{48}{143}\right)\!=\!\left(\!\frac{5}{\Gamma\left(\frac{1}{2}\!+\!1\right)}\!+\!\frac{4}{\Gamma\left(\frac{1}{2}\!\right)}\!\right)\!\frac{1}{50}\!+\!\left(\frac{48}{143}\right)\\ &=\left(\frac{10}{\sqrt{\pi}}\!+\!\frac{4}{\sqrt{\pi}}\right)\!\frac{1}{50}\!+\!\left(\frac{48}{143}\right)\!=\!\left(\frac{14}{50\sqrt{\pi}}\right)\!+\!\left(\frac{48}{143}\right)\!<1. \end{split}$$

Similarly,

$$\begin{split} &\sigma_2 = \left(\frac{1}{\Gamma(m+1)} + \frac{2(\theta+\vartheta)}{\Gamma(m+1)} + \frac{2(\theta^2+\theta\vartheta)}{\theta^2\Gamma(m)}\right) \Re_k + \frac{(\bar{\theta}+\bar{\vartheta})\left(\Re_{\bar{b}_1} + \Re_{\bar{b}_2}\right)}{\bar{\theta}^2} \\ &= \left(\frac{1}{\Gamma(m+1)} + \frac{4}{\Gamma(m+1)} + \frac{4}{\Gamma(m)}\right) \frac{1}{49} + 2\left(\frac{1}{11} + \frac{1}{13}\right) \\ &= \left(\frac{5}{\Gamma(m+1)} + \frac{4}{\Gamma(m)}\right) \frac{1}{49} + \left(\frac{48}{143}\right) = \left(\frac{5}{\Gamma\left(\frac{1}{2}+1\right)} + \frac{4}{\Gamma\left(\frac{1}{2}\right)}\right) \frac{1}{49} + \left(\frac{48}{143}\right) \\ &= \left(\frac{10}{\sqrt{\pi}} + \frac{4}{\sqrt{\pi}}\right) \frac{1}{49} + \left(\frac{48}{143}\right) = \left(\frac{14}{49\sqrt{\pi}}\right) + \left(\frac{48}{143}\right) < 1. \end{split}$$

Consequently, it can be asserted that the coupled system denoted as (5.1) possesses a singular solution. Furthermore, the condition stipulated in Theorem 2, expressed as  $(1 - \Re_{\theta \vartheta})(1 - \Re_{\bar{\theta} \vartheta}) - \Re_{m\theta \vartheta} \Re_{n\bar{\theta} \bar{\vartheta}} \neq 0$ , is also satisfied.

Consequently, the coupled system (5.1) demonstrates both Ulam-Hyers stability and generalized Ulam-Hyers stability.

# **CONCLUSION**

We investigate Ulam-Hyers and generalized Ulam-Hyers stability for fractional integro-differential coupled systems with integral boundary conditions, an area where existing research has focused mainly on simpler, isolated equations rather than the coupled systems that appear more frequently in practice. Our approach differs by analyzing how these coupled systems behave when subjected to small perturbations, using fixed-point theorems and carefully constructed function spaces to establish both solution existence and uniqueness.

What makes this work particularly valuable is its relevance to real applications in engineering, finance, and physics, where model stability directly impacts whether computational results can be trusted for decision-making. Our quantitative analysis shows that when initial conditions vary slightly, the resulting solution changes remain bounded and predictable, exactly the kind of stability guarantee practitioners need. Beyond these immediate results, the mathematical framework we've developed opens doors to studying more complex systems, including those with neutral delays, impulsive effects, or various inclusion properties that arise when modeling systems with both finite and infinite time delays. Rather than simply filling theoretical gaps, this work provides concrete tools that researchers can apply when analyzing stability in increasingly sophisticated differential systems.

#### **ACKNOWLEDGMENTS**

The authors thank the reviewers for their valuable comments and suggestions that improved this manuscript.

All authors have read and approved the final manuscript.

#### **AUTHORSHIP CONTRIBUTIONS**

Brijendra Kumar Chaurasiya: Investigation, Methodology, Validation, Writing-original draft, Writingreview and editing. Avadhesh Kumar: Conceptualization, Investigation, Methodology, Supervision, Validation, Writing-original draft, Writing-review and editing.

#### **DATA AVAILABILITY STATEMENT**

The authors confirm that the data that supports the findings of this study are available within the article. Raw data that support the finding of this study are available from the corresponding author, upon reasonable request.

# **CONFLICT OF INTEREST**

The author declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

#### **ETHICS**

There are no ethical issues with the publication of this manuscript.

# STATEMENT ON THE USE OF ARTIFICIAL INTELLIGENCE

Artificial intelligence was not used in the preparation of the article.

#### **REFERENCES**

- [1] Chalishajar D, Kumar A. Existence, uniqueness and Ulam's stability of solutions for a coupled system of fractional differential equations with integral boundary conditions. Mathematics 2018;6:96.

  [CrossRef]
- [2] Kumar A, Kumar A, Vats RK, Kumar P. Approximate controllability of neutral delay integrodifferential inclusion of order α∈(1,2) with non-instantaneous impulses. Evol Equ Control Theory 2022;11:1635–1654. [CrossRef]
- [3] Yang Q, Chen D. Fractional calculus in image processing: a review. Fract Calc Appl Anal 2016;19:1222–1249. [CrossRef]
- [4] Magin RL. Fractional calculus in bioengineering. Redding (CA): Begell House; 2006.
- [5] Kilbas AA, Srivastava HM, Trujillo JJ. Theory and applications of fractional differential equations. Vol. 204. New York: Elsevier Science; 2006.
- [6] Miller KS, Ross B. An introduction to fractional calculus and fractional differential equations. New York: Wiley Interscience; 1993.
- [7] Hilfer R. Applications of fractional calculus in physics. Singapore: World Scientific; 2000. [CrossRef]
- [8] Khan A, Shah K, Li Y, Khan TS. Ulam type stability for a coupled system of boundary value problems of nonlinear fractional differential equations. J Funct Spaces 2017;2017:1–8. [CrossRef]
- [9] Ahmad B, Nieto JJ. Existence results for nonlinear boundary value problems of fractional integro-differential equations with integral boundary conditions. Bound Value Probl 2009;2009:1–11. [CrossRef]
- [10] Jafari H, Mehdinejadiani B, Baleanu D. Fractional calculus for modeling unconfined groundwater: applications in engineering, life and social sciences. Mathematics 2019;7:119–138. [CrossRef]
- [11] Oldham KB. Fractional differential equations in electrochemistry. Adv Eng Softw 2010;41:9–12. [CrossRef]
- [12] Jannelli A. Numerical solutions of fractional differential equations arising in engineering sciences. Mathematics 2020;8:215. [CrossRef]
- [13] Ionescu C, Lopes A, Copot D, Machado JT, Bates JH. The role of fractional calculus in modeling biological phenomena: a review. Commun Nonlinear Sci Numer Simul 2017;51:141–159. [CrossRef]

- [14] Ulam SM. A collection of the mathematical problems. New York: Interscience; 1960.
- [15] Hyers DH. On the stability of the linear functional equation. Proc Natl Acad Sci U S A 1941;27:222–224. [CrossRef]
- [16] Jung SM. Hyers-Ulam stability of linear differential equations of the first order, II. Appl Math Lett 2006;19:854–858. [CrossRef]
- [17] Wang J, Feckan M. A general class of impulsive evolution equations. Topol Methods Nonlinear Anal 2015;46:915–933. [CrossRef]
- [18] Muslim M, Kumar A, Feckan M. Existence, uniqueness and stability of solutions to second-order non-linear differential equations with non-instantaneous impulses. J King Saud Univ Sci 2016;28:1–9.
- [19] Rus IA. Ulam stabilities of ordinary differential equations in a Banach space. Carpath J Math 2010;26:103–107.
- [20] Ahmad B, Sivasundaram S. Existence and uniqueness results for nonlinear boundary value problems of fractional differential equations with separated boundary conditions. Commun Appl Anal 2009;13:121. [CrossRef]
- [21] Verma P, Kumar M. Analytical solution with existence and uniqueness conditions of non-linear initial value multi-order fractional differential equations using Caputo derivative. Eng Comput 2022;38:661–678. [CrossRef]
- [22] Van Bockstal K, Zaky MA, Hendy AS. On the existence and uniqueness of solutions to a nonlinear variable order time-fractional reaction-diffusion equation with delay. Commun Nonlinear Sci Numer Simul 2022;115:106755. [CrossRef]
- [23] Shafqat R, Niazi AUK, Jeelani MB, Alharthi NH. Existence and uniqueness of mild solution where α∈(1,2) for fuzzy fractional evolution equations with uncertainty. Fractal Fract 2022;6:65.
- [24] Sweis H, Shawagfeh N, Abu Arqub O. Fractional crossover delay differential equations of Mittag-Leffler kernel: existence, uniqueness, and numerical solutions using the Galerkin algorithm based on shifted Legendre polynomials. Results Phys 2022;41:105891. [CrossRef]
- [25] Younis M, Singh D. On the existence of the solution of Hammerstein integral equations and fractional differential equations. J Appl Math Comput 2022;68:1087–1105. [CrossRef]
- [26] Telli B, Souid MS, Alzabut J, Khan H. Existence and uniqueness theorems for a variable order fractional differential equation with delay. Axioms 2023;12:339. [CrossRef]
- [27] Mahdy AMS. Stability, existence, and uniqueness for solving fractional glioblastoma multiforme using a Caputo-Fabrizio derivative. Math Methods Appl Sci 2023;46:1–15.

- [28] Mehmood N, Abbas A, Akgül A, Abdeljawad T, Alqudah MA. Existence and stability results for coupled system of fractional differential equations involving AB-Caputo derivative. Fractals 2023;31:2340023. [CrossRef]
- [29] Zou J, Luo D, Li M. The existence and averaging principle for stochastic fractional differential equations with impulses. Math Methods Appl Sci 2023;46:6857–6874. [CrossRef]
- [30] Khan H, Alzabut J, Baleanu D, Alobaidi G, Rehman MU. Existence of solutions and a numerical scheme for a generalized hybrid class of n-coupled modified ABC-fractional differential equations with an application. Math Methods Appl Sci 2023;46:1–15.
- [31] Toprakseven S, Bengi R, Zeren Y. Lyapunov-type inequality for an anti-periodic fractional boundary value problem of the Riesz-Caputo derivative. Sigma J Eng Nat Sci 2023;41:807–811. [CrossRef]
- [32] Adıgüzel H. A note on the asymptotic behavior of fractional differential equations. Sigma J Eng Nat Sci 2020;38:1061–1067.
- [33] Koşunalp HY, Gülsu M. Operational matrix for multi-order fractional differential equations with Hermite polynomials. Sigma J Eng Nat Sci 2024;42:1050–1057. [CrossRef]
- [34] Adıgüzel H. A note on asymptotic behavior of fractional differential equations. Sigma J Eng Nat Sci 2020;38:1061–1067.
- Jan R, Yüzbaşı Ş, Attaullah A, Jawad M, Jan A. Fractional derivative analysis of asthma with the effect of environmental factors. Sigma J Eng Nat Sci 2024;42:177–188. [CrossRef]
- [36] Sanatee AG, Rathour L, Mishra VN, Dewangan V. Some fixed point theorems in regular modular metric spaces and application to Caratheodory's type anti-periodic boundary value problem. J Anal 2023;31:619–632. [CrossRef]
- [37] Shahi P, Rathour L, Mishra VN. Expansive fixed point theorems for tri-simulation functions. J Eng Exact Sci 2022;8:14303–01e. [CrossRef]
- [38] Pathak VK, Mishra LN, Mishra VN, Baleanu D. On the solvability of mixed-type fractional-order non-linear functional integral equations in the Banach space C(l). Fractal Fract 2022;6:744. [CrossRef]
- [39] Metwali MM, Mishra VN. On the measure of non-compactness in Lp(R+) and applications to a product of s-integral equations. Turk J Math 2023;47:372–386. [CrossRef]
- [40] Bouzalmat I. Stochastic modeling of COVID-19 dynamics with emphasis on asymptomatic and isolated populations. Proc Int Conf Math Model Mod Appl 2024;1:1–12.
- 41] Paul SK, Mishra LN, Mishra VN, Baleanu D. An effective method for solving nonlinear integral equations involving the Riemann-Liouville fractional operator. AIMS Math 2023;8:17448–17469. [CrossRef]

- [42] Paul SK, Mishra LN, Mishra VN, Baleanu D. Analysis of mixed-type nonlinear Volterra-Fredholm integral equations involving the Erdélyi-Kober fractional operator. J King Saud Univ Sci 2023;35:102949.

  [CrossRef]
- [43] Bhat IA, Mishra LN, Mishra VN, Tunc C, Tunc O. Precision and efficiency of an interpolation approach to weakly singular integral equations. Int J Numer Methods Heat Fluid Flow 2024;34:1479–1499. [CrossRef]
- [44] Kumar K, Rathour L, Sharma MK, Mishra VN. Fixed point approximation for Suzuki generalized nonexpansive mapping using  $B(\delta,\mu)$  condition. Appl Math 2024;13:215–227. [CrossRef]
- [45] Liu X, Zhou M, Ansari AH, Chakrabarti K, Abbas M, Rathour L. Coupled fixed point theorems with rational type contractive condition via c-class functions and inverse ck-class functions. Symmetry 2022;14:1663. [CrossRef]
- [46] Iqbal J, Mishra VN, Mir WA, Dar AH, Ishtyak M, Rathour L. Generalized resolvent operator involving  $G(\cdot,\cdot)$ -co-monotone mapping for solving generalized

- variational inclusion problem. Georgian Math J 2022;29:533–542. [CrossRef]
- [47] Maheswari U, Ravichandran M, Anbarasan A, Rathour L, Mishra VN. Some results on coupled fixed point on complex partial b-metric space. Ganita 2021;71:17–27.
- [48] Deng J, Liu X, Sun Y, Rathour L. On almost type  $\alpha$ -FZ-weak contraction in metric spaces. Open J Appl Sci 2022;12:528–540. [CrossRef]
- [49] Deng J, Liu XL, Sun Y, Rathour L. Some best proximity point results of several  $\alpha$ - $\psi$  interpolative proximal contractions. Nonlinear Funct Anal Appl 2022;27:533–551.
- [50] Mishra AK, Rathour Padmawati L, Mishra VN. Some fixed point theorem using generalized cone metric spaces with Banach algebra. High Tech Lett 2023;29:153–162.
- [51] Alam M, Zada A, Abdeljawad T. Stability analysis of an implicit fractional integro-differential equation via integral boundary conditions. Alex Eng J 2024;87:501–514. [CrossRef]